



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed



for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display Abstract



Show

20



Sort by



Send to



About Entrez

Text Version

All: 1

Review: 0



Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Mol Chem Neuropathol. 1998 Jun-Aug;34(2-3):121-32.

Related Articles. Links

## Extensive degeneration of catecholaminergic neurons to scrapie agent 87V in the brains of IM mice.

Yun SW, Choi EK, Ju WK, Ahn MS, Carp RI, Wisniewski HM, Kim YS.

Institute of Environment and Life Science, College of Medicine, Hallym University, Chunchon, Korea.

Scrapie is a degenerative disease of the central nervous system of sheep and goats. The causative agent has been passaged to a number of laboratory species, including mice and hamster. Amyloid plaque formation and vacuolation, the signs of senile dementia, are found in the brains of mice infected with 87V scrapie agent. Dopamine (DA) and norepinephrine (NE) concentrations in the brains of scrapie-infected mice were measured with high-performance liquid chromatography-electrochemical detector (HPLC-ECD). A significant decrease in NE level was exhibited in all regions tested, whereas the level of DA decreased significantly only in cerebral cortex. Immunohistochemistry was used to examine immunoreactive catecholamine neurons in substantia nigra and locus ceruleus using antisera against tyrosine hydroxylase (TH). The population of TH-immunoreactive neurons in the substantia nigra and locus ceruleus were significantly decreased in scrapie-infected mice compared to controls. These data suggest that both the noradrenergic and dopaminergic system are sensitive to the action of scrapie agent 87V and that changes in the catecholamine levels in the brains of scrapie-infected mice may contribute to some of the clinical symptoms of the diseases, such as ataxia and apraxia.

PMID: 10327412 [PubMed - indexed for MEDLINE]

Display Abstract



Show

20



Sort by



Send to



[Write to the Help Desk](#)

[NCBI | NLM | NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Aug 16 2005 14:03:09